

Hip-Spine

Back to ignoring it all over again

Jeff Barry, MD
Associate Clinical Professor of Orthopaedics
Arthroplasty Fellowship Director
UCSF Department of Orthopaedic Surgery





Disclosures

- •Consulting:
 - Smith & Nephew
 - Onkos
 - J&J MedTech
 - Lineage Medical
- •Institutional Fellowship grants: S&N, AAHKS




Outline

 Understand Pelvic Position and the relationship to Acetabular Position

Barry Algorithm for Cup Positioning



Hip-Spine Recently





Don't Be This Guy Today










A Way Without Classifications or Measurements or Robots or Sit to Stand Spine Films









Why is Spine a Topic of Every Meeting



•Alternative Topics

- **C**Exactech®
- Poly wear (solved)
- CoCr issues (solved)
- Anterior vs Posterior hips (solved)
- Infection (never gonna solve)



Posterior hips do just as well!!!




- Let's talk about the spine!!
 - Admittedly has led to improved biomechanical understanding





The Stable (Native) Hip Joint

- Enarthrodial ball and socket joint
- Inherently stable
 - Bony congruity
 - Labrum
 - Capsulo-ligamentous complex











Human Pelvis and Hip Joint

 Evolved to allow upright bipedal ambulation and still be able to birth large brains










Pelvic Terms

- Measurements inherent to pelvic anatomy
 - Pelvic incidence

- Dynamic / positional / functional measures
 - Pelvic tilt, sacral slope, lumbar lordosis
 - Acetabular anteversion and inclination










Pelvic Incidence

- Fixed in maturity Increases during development
- Measure of *individual* pelvic anatomy














Pelvic Incidence (fixed) Sets Lordosis

■ Sets the base of the spine









Work What Your Momma Gave You








Pelvic Tilt – <u>Positional</u> Parameter

- Spine measurement
 - Normal ~13deg

- Arthroplasty measurement –ASIS to symphysis
 - Look at anterior pelvic plane
 - ~zero normal




PELVIC TILT CAN BE APPROXIMATED ON AP IMAGE

MORE INLET = ANTERIOR TILT

TILTED

PELVIS


MORE OUTLET = POSTERIOR TILT





The pelvis was meant to tilt













The pelvis was meant to tilt

- Standing to sitting: PT increases (roll back)
 - Opens up cup to allow legs to flex up without impingement
- Lying down PT decreases (roll forward)
 - Coverage against anterior instability
 - Important in DA surgery positioning!

 Changes in PT increase safe functional range of motion without impingement in native hip










Effect of Stiff Spine

- Stuck Sitting pelvis already retroverted can't roll forward when standing and can get impingement in extension (anterior instability)
- Stuck Standing pelvis neutral and can't roll on sitting risk anterior impingement (posterior instability









Connection of Tilt and Acetabular Position






THE DEFINITION AND MEASUREMENT OF ACETABULAR ORIENTATION

D. W. MURRAY

- Typically anteversion refers to "operative" anteversion
- How much you raise your hand from longitudinal axis












Operative Anteversion

- Hard to measure radiographically so surrogates are used (and abused...)
 - Anteinclination
 - Estimates based on major and minor diameters on an AP
 - Ackland
 - Projection anteversion
 - Anatomic anteversion (CT)
 - Lateral anteversion
 - Acetabular tilt

•All are positional!



















Acetabular Position Changes with Pelvic Tilt








Pelvic/Spine Summary

- Pelvis and spine moves depending on functional position
- Stiff spines can be stuck in various tilted positions
- Acetabular position is a function of pelvic position and tilt

■ What isn't here... clinical implications!!!!








Dislocations after Total Hip-Replacement Arthroplasties*

BY GEORGE E. LEWINNEK, M.D.†, JACK L. LEWIS, PH.D.‡, RICHARD TARR, M.S.‡, CLINTON L. COMPERE, M.D.†,
AND JERALD R. ZIMMERMAN, B.S.‡, CHICAGO, ILLINOIS

From the Northwestern Memorial Hospital, Chicago

- 9 dislocations
- Defined a safe zone
 - 40 +/- 10° inclination
 - 15 +/- 10° anteversion
- 1.5% dislocation rate in safe zone
- 6% dislocation rate out of safe zone
- Orthopaedic gospel set by 3 cases you should ignore and 6 cases you should listen to
- NO CONCEPT OF PELVIC POSITIONING OR STANDARDIZATION ON RADIOGRAPHS!!!!!





Summary of 1882 Papers on Spine and Hip Clinical Outcomes

■Bad spine = more risk for problems after THA










How does this impact what I should do?





Who's At Risk.... (taken from AAHKS FOCAL lecture)



J. Langston. J. Pierrepont,

From Mclbourne

Orthopedic Group.

Windsor, Australia

Y. Gu, A. Shimmin

motion causing unfavourable ori the acetabular component in pat undergoing total hip arthroplasty

It is important to consider sagittal pelvic rotation when introducing the acetabular component at total hip arthroplasty (THA). The purpose of this study was to identify patients who are at risk of unfavourable pelvic mobility, which could result in poor outcomes after THA.

A consecutive series of 4042 patients undergoing THA had lateral functional radiographs and a low-dose CT scan to measure supine pelvic tilt, pelvic incidence, standing pelvic tilt, flexedseated pelvic tilt, standing lumbar lordotic angle, flexed-seated lumbar lordotic angle, and lumbar flexion. Changes in pelvic tilt from supine-to-standing positions and supine-to-flexedseated positions were determined. A change in pelvic tilt of 13° between positions was

- Standing Posterior PT > 10°
- 2. PT Change Supine → Standing ≤ -13°
- 3. PT Change Stand → Seated ≥ 20°

SHOW YOUR PRODUCTION AND PARTY \$JJ-2017-1995/RT 52:00

Boow Joint J

VOL. 100-8, No. 7, BULY 2015

mately 31. If the pelvis tilts posteriorly, there is however, dislocations continued. 58

established that a change in sagittal pervicibil of commended a safe zone for inclination of 10° will change the version of the acetabulum by 40° ± 10° and for anteversion of 15° ± 10°. approximately 7s and its inclination by approxi- Despite placing components in these positions,

Risk factors for increased sagitta motion causing unfavourable original property of the control of the control

CONFUSED!

The Journal of Arthroplasty 32 (2017) 2310-1917

Contents lists available at ScienceDirect

The Journal of Arthroplasty

al homepage: www.arthroptestyjournel.prg



Total Hip Arthroplasty in the Spinal Deformity Population: Does Degree of Sagittal Deformity Affect Rates of Safe Zone Placement, nstability, or Revision?



Edward M. DelSole, MD *, Jonathan M. Vigdorchik, MD, Ran Schwarzkopf, MD, MSc. Thomas J. Errico, MD, Aaron J. Buckland, MBBS, FRACS

Department of Orthopaedic Suppry. The Hospital for John Diseases, NAI Language Medical Center, New York, New York

ARTICLE INFO

ABSTRACT

* Department of Biomechanics, Hospital for Special Surgery, New York, NY

Adult Speciestrochini and Joint Replacement District. Department of Orthopardit Surgery, Nanpini for Special Surgery, New York, NY

fluckground: Sitting ratiographs have been used as a pre-operative tool to plan patient-specific total hip-

4. Lumbar Flexion ≤ 20° **Have Significantly Higher Dislocation Rates** 5. Advanced Age >75 Yrs Especially In Females

radiographic acetabular "safe zone" that has been widely used since 1978 does not guarantee decreased risk of hip dislocation [3,4]. This can be explained by the fact that a standard anteroposterior

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may suckade receipt of payment, either direct or indirect, institutional support, or association with an entity in the bursedical field which may be perceived to have potential conflict of interest with this mark. For full disclosure seasoneers neter to https://doi.org/10.1036/Sants.303733.005 * Repoint opparate: Christina I. Esposito, Phil), Department of Biomechanics Hospital for Special Surgery, 535 East 70th Street, New York, NY 10021.

herom, Fisher year 100 toll (A. 2005, 2017) 22 (2015) 0883-5403(G-2017 Essevint Inc. All rights reserved.

which aid in predicting an ideal, patient-specific THA component position that would improve stability and limit-bearing surface wear. How parients achieve sitting position through the lumbarpelvic-femoral complex continues to be an area of interest given the impact on functional acetabular component position following THA [3.3,7], and arthroplasty surgeons are becoming increasingly aware that previous spine surgery may be associated with poor outcomes after THA [5,9].

Sitting position can be achieved through a combination of lumbar spine segmental motions and hip motion. Patients with stiff spines have a limited change in pelvic tilt from standing to sitting 6. Severe Sagittal Spinal **Deformity (Pelvic** Incidence - Standing LL Mismatch ≥ 20°) Have **Dislocation Rate Of 8%**

sopelvic tilt (SPT) and, as a consequence, changes the functional position of the acetabulum. This has a direct impact on the


One or more of the authors of this paper have discissed potential or partment coefficis of extensit, which may include receipt of payment, either direct or indirect, nutifulional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to http://doi.org/10.10163.arch.2016.12.009.

stigation was performed at NYU Hospital for Joint Diseases, New York, NY. Reprint requests: Edward M. DelSole, MD, Department of Orthogoedic Surgery The Huspital for Joint Discuses. NYU Langone Medical Center, 301 East 17th Street, luite 1402, New York, NY 10003.

eter (18x 8x2.com) tit 1016/15arth 2016.12.076 0\$83-5403/s: 2016 Eurwier Inc. All rights reserved. and kyphotic curvature of the spiral column, resulting in abnorma pelvic or lower extremity posturing in an attempt to regain balance [10]. The role of the pelvis in compensating for SSD is essential; as a patient's overall sagittal balance tends toward kyphosis, patients increase pelysc tilt in an effort to maintain upright posture and forward gaze [1],32]. This pelvic compensation may cause unexpected changes in cup position during transition to a standing position, leading to instability of the hip prosthesis [13].

In this study, we sought to investigate the acetabular cup position, dynamic changes in cup position, and rate of instability episodes in patients with SSD who have undergone THA. We evaluated the relationships between cup position in both the











Barry "Algorithmic" Approach to Spine Patient

- 1. Preop standing AP pelvis (define functional plane)
- 2. Antevert cup from the functional plane (match the tilt)
 - 20-25 deg for anterior approach
 - 25-35 deg for posterior approach








Barry "Algorithmic" Approach to Spine Patient

*** Exception in primary <u>posterior</u> hips:

• Long segment fusion to pelvis (>3 levels) consider DM

• Uncoordinated frail grandma scares me way more than yoga instructor













I'm going to say the same thing but make it more confusing

C.S. Sicat et al. / The Journal of Arthroplasty 37 (2022) S540-S545

Group	Classification Normal Alignment Normal Mobility	Explanation of Pathology	Treatment Recommendation Target 20°-25° anteversion from normal anatomy (to the APP)	
1A		Normal Anatomy and Normal Mobility (APP similar to FPP)		
1B	Normal Alignment Stiff Spine (<10° change in Sacral Slope from stand to sit)	Stiff spine, needs more anteversion relative to the APP which is similar to the FPP	Target 30° anteversion on the standing AP Pelvis (to the FPP). Note the higher target due to the stiff spine to protect from a posterior dislocation.	
2A	Flatback deformity (PI-LL>10°) Normal Mobility	Be aware the FPP is different than the APP (posterior pelvic tilt from the spinal deformity causes more functional cup anteversion)	Target 25°-30° anteversion on the <u>standing</u> AP Pelvis (to the FPP)	
2B*	Flatback deformity (PI-LL>10°) Stiff Spine (<10°change in Sacral Slope from stand to sit)	FPP is different than the APP and a stiff spine, so you need more anteversion, but be aware of the reference because the spinal deformity will cause more functional cup anteversion	Target 30° anteversion on the standing AP Pelvis (to the FPP). Note the higher target due to the stiff spine to protect from a posterior dislocation.	


- 1. <u>Preop standing AP pelvis</u> (Functional position)
- 2. Antevert from the functional position
 - 20-25deg for DA;
 - 25-35deg for posterior

Classification made for posterior approach

Barry Algorithm fits into every category! (25-35deg anteversion for posterior hips to standing AP Pelvis)



Fig. 1. Explanation of Hip-Spine Classification and corresponding treatment recommendations. FPP, functional pelvic plane; APP, anatomic pelvic plane.



Jeff, you said there was no measurements. 20-25 or 25-35 how do I calculate or get there? What gives?



The Journal of Arthroplasty Vol. 19 No. 3 2004

Math to Ignore

A Simplified Method to Determine Acetabular Cup Anteversion From Plain Radiographs

Karl-Heinz Widmer, MD, Dipl Ing (TU)




Table 1. S/TL-Ratio Versus Radiographic Cup Anteversion on a Plain Radiograph Centered on the Symphysis for 40°, 45°, 50°, and 55° Cup Inclination*

	Radiographic Cup Anteversion Cup Inclination				
S/TL	40°	45°	50°	55°	
0.00	3.5°	3.9°	4.2°	4.5°	
0.04	4.6°	5.0°	5.2°	5.5°	
0.07	5.6°	6.0°	6.3°	6.6°	
0.11	6.7°	7.1°	7.4°	7.7°	
0.14	7.8°	8.1°	8.4°	8.7°	
0.17	8.8°	9.2°	9.5°	9.8°	
0.20	9.9°	10.3°	10.6°	10.8°	
0.23	11.0°	11.3°	11.6°	11.9°	
0.26	12.0°	12.4°	12.7°	13.0°	
0.29	13.1°	13.5°	13.8°	14.1°	
0.31	14.2°	14.6°	14.9°	15.1°	
0.34	15.3°	15.7°	15.9°	16.2°	
0.36	16.3°	16.7°	17.0°	17.3°	
0.39	17.4°	17.8°	18.1°	18.4°	
0.41	18.5°	18.9°	19.2°	19.5°	
0.43	19.6°	20.0°	20.3°	20.6°	
0.46	20.7°	21.1°	21.4°	21.7°	
0.48	21.9°	22.2°	22.5°	22.8°	
0.50	23.0°	23.4°	23.7°	23.9°	
0.52	24.1°	24.5°	24.8°	25.1°	
0.54	25.2°	25.6°	25.9°	26.2°	
0.56	26.4°	26.8°	27.1°	27.4°	
0.58	27.6°	27.9°	28.2°	28.5°	
0.60	28.7°	29.1°	29.4°	29.7°	
0.62	29.9°	30.3°	30.6°	30.9°	
0.63	31.1°	31.5°	31.8°	32.0°	
0.65	32.3°	32.7°	33.0°	33.3°	
0.67	33.5°	33.9°	34.2°	34.5°	
0.68	34.7°	35.1°	35.4°	35.7°	
0.70	36.0°	36.4°	36.7°	37.0°	
0.71	37.3°	37.6°	37.9°	38.2°	
0.73	38.5°	38.9°	39.2°	39.5°	
0.74	39.9°	40.2°	40.5°	40.8°	

Anteversion from Projection

SET FUNCTIONAL PELVIC PLANE
(MATCH STANDING AP INLET/OUTLET APPEARANCE)

Anterior Approach


Solid to Open 50/50 - 1:1

= ~23deg anteversion




Solid to Open 33/66 or 1:2

= ~32deg anteversion














Why This Works!!!!

Instability is not JUST cup position

- Component(S) position femoral version matters as much as cup
- Impingement source
 - USUALLY SOFT TISSUE OR TROCH ON PELVIS NOT NECK ON CUP!!!
- Anatomy restoration
- Other (spine, muscle quality, abductor insufficiency, FNF or AVN, dementia)
- Approach





Approach and Instability

- ■DA is not DA is not DA....
 - -PRESERVE THE POSTERIOR CAPSULE AND ITS HARD TO COME OUT THE BACK!!!
 - DON'T do inferior episiotomy
 - -Repair anterior capsule
 - –Minimize femoral releases (table is your best instrument)
 - -Recreate native offset
 - Anterior positions of instability are unnatural



Approach and Instability

- Posterior approach
 - Template to restore anatomy
 - can augment with robot or navigation if not confident
 - -Trial to determine instability sources
 - Troch vs hypertrophic capsule vs thigh on belly vs implant impingement (RARE)
 - Repair posterior capsule when able
 - -Intraoperative xray MATCHING STADNING before finalizing cup






DA Advantage = Fluoro Advantage!!!

- Match fluoro plane to standing view (add ~10deg outlet) (Roettes JoA 2018)
- Center over the cup to avoid parallax in old carm
 - Square intensifier = vascular = no parallax
 - Round intensifier = parallax susceptible
 - (test with metal rod if not sure)


50/50 Open/Closed in the functional positional



NOTE** if not a vascular style digital carm center your carm over the area of interest (ie cup during insertion)









Conclusions

- Acetabular position is a dynamic measure based on pelvic position
- Standing AP Pelvis tells you all you need to know about the spine
 - FUNCTIONAL PELVIC CUP POSITIONING (and anterior approach) will keep you safe
- Intraop Targets based on the FUNCTIONAL POSITION (Standing AP Pelvis Match)
 - Anterior 50/50 cup
 - Posterior 66/33 cup
- Anterior surgeons go back to ignoring spine
- Posterior surgeons go back to ignoring spine (except when you see lots of hardware)
 - DM judiciously





